

VOLUME 38, NUMBER 4
APRIL 2021

ISSN 0189 - 160X

WAJM

WEST AFRICAN JOURNAL OF MEDICINE

ORIGINALITY AND EXCELLENCE IN MEDICINE AND SURGERY



OFFICIAL PUBLICATION OF
THE WEST AFRICAN COLLEGE OF PHYSICIANS AND
WEST AFRICAN COLLEGE OF SURGEONS



www.wajmed.org



TABLE OF CONTENTS

GENERAL INFORMATION	IC
INFORMATION FOR AUTHORS	1F
EDITORIAL NOTES	299
ORIGINAL ARTICLES	
HIV-Associated Nephropathy among Children with Renal Disease in Port Harcourt, Nigeria	307
T. A. Uchenwa, I. C. Anochie	
Dental Trauma in Adult and Elderly Nigerians: A National Survey	313
E. C. Otoh, O. O. Taiwo, O. A. Adeleke, O. J. Majekodunmi, S. O. Ajike	
Knowledge, Attitude and Practice on Covid-19 among Clinical Healthcare Workers in Bingham University Teaching Hospital (BHUTH) Jos, Plateau State, Nigeria	321
M. Shehu, H. Shehu, O. Momodu, O. Abraham, E. E. Eseigbe	
Comparison of the Diagnostic Relevance of Albumin Creatinine Ratio Versus Cystatin C in Assessment of Cardiovascular Complication in Type 2 Diabetics	328
B. K. Myke-Mbata, S. C. Meludu, D. O. Ochalefu, B. C. Basil	
Self-perceived Burden on Caregivers, Anxiety and Depression among Chronic Kidney Disease Patients in Southern Nigeria	335
O. A. Adejumo, E. I. Okaka, A. A. Akinbodewa, O. I. Iyawe, I. R. Edeki, O. S. Abolarin	
Frequency of Osteoporosis in Black Nigerian Women Aged 50 and above with Degenerative Musculoskeletal Diseases and Fractures	342
O. A. Adewole, S. O. Idowu, M. O. Shoga, M. O. Kayode, O. O. Adelowo	
Knowledge, Attitudes, and Practices towards COVID-19 Transmission and Preventive Measures among Residents of Nigeria: A Population-Based Survey through Social Media	347
A. T. Akafa, A. Amos , A. Okeke and A. C. Oreh	
School Health Instructions in Primary Schools - A Study of Gwagwalada Area Council, Federal Capital Territory Nigeria	359
U. A. Sanni, U. M. Offiong, E. A. Anigilaje, K. I. Airede	
Plasma L-Arginine in Sickle Cell Anaemia Patients in Crises and its Correlation with Markers of Severity of Disease	366
O. W. Aworanti, T. S. Akingbola, A. Adeomi, A. E. Alagbe, A. O. Salako	
Meniscus Sign: A Test for the Confirmation of Correct Placement of Epidural Catheter	374
H. O. Idehen, G. Agbonkhese	
Clinicopathological Pattern and Management of Primary Lung Cancer in Ilorin, Nigeria	380
P. O. Adeoye, O. O. Desalu, C. K. P. Ofoegbu, A. E. Fawibe, A. K. Salami, R. O. Akanbi, C. M. Opeyemi, O. O. K. Ibrahim, O. B. Ojuawo, A. O. Aladesanmi, I. Olaoye	
CASE REPORTS	
Protein C Deficiency in a Patient with Anomalous Hemiazygous Vein and Portal Vein Thrombosis	387
F. A. Fasola, A. Akere, C. U. Akunwata, C. Onyejelam, M. A. Osundina	
Aplasia Cutis Congenita: A Case Report	391
O. J. Ugowe, S. A. Balogun and E. A. Adejuyigbe	
Favourable Outcome of Severe Lassa Fever Following Early Diagnosis and Treatment: A Case Report	395
J. A. Onuh, A. E. Uloko	
CLINICAL ARTICLE	
Comparison of Short Course Versus Long Course Antibiotic Prophylaxis for Caesarean Section: A Randomised Controlled Trial	398
A. C. Ezeike, C. O. Agboghoroma, E. R. Efetie, K. W. Durojaiye	
INDEX TO VOLUME 38, NO. 4, 2021	
Author Index	405
Subject Index	406



ORIGINAL ARTICLE

Self-perceived Burden on Caregivers, Anxiety and Depression among Chronic Kidney Disease Patients in Southern Nigeria

Fardeau auto-perçu des soignants, anxiété et dépression chez les patients atteints d'insuffisance rénale chronique dans le sud du Nigéria

¹O. A. Adejumo, ²E. I. Okaka, ¹A. A. Akinbodewa, ²O. I. Iyawe, ²I. R. Edeki, ¹O. S. Abolarin

ABSTRACT

Introduction: Psychosocial problems are common in chronic kidney disease (CKD) patients and are associated with poor overall outcomes; however, their evaluation has not received the deserved attention. This study assessed self-perceived burden on caregivers, psychosocial wellbeing, anxiety and depression among CKD patients, and the impact on their quality of life (QoL) in two hospitals in Nigeria.

Methods: This was a cross-sectional study. Self-perceived burden on caregivers, QoL, anxiety, and depression of the patients, were assessed using the Cousineau questionnaire, modified SF-12 questionnaire, Hospital Anxiety and Depression Scale, respectively.

Results: There were 141 participants in the study comprising 50 haemodialysis (HD) patients, 41 pre-dialysis CKD patients and 50 controls. The male:female ratio for CKD patients and controls were 1.1 and 1.4 respectively. Prevalence of depressive and anxiety symptoms in the CKD patients were 46.2% and 33.0% respectively. The QoL score was significantly higher in the control group compared to the CKD patients ($p = <0.001$) and significantly lower in the HD patients compared to the pre-dialysis CKD patients ($p = <0.001$). Self-perceived burden score was significantly higher in the HD group compared to the pre-dialysis CKD group ($p = <0.001$). There was a negative correlation between QoL score and self-perceived burden, hospital anxiety score and hospital depression score ($p = <0.001$).

Conclusion: Psychological disorders and self perceived burden are highly prevalent among CKD patients. They have negative impact on their QoL, hence regular evaluation and management of these disorders should be incorporated into the care of CKD patients. **WAJM 2021; 38 (4): 335-341**

Keywords: Anxiety, Depression, Quality of Life, Burden on caregivers, chronic kidney disease

ABSTRAIT

Introduction: Les problèmes psychosociaux sont commun chez les patients atteints d'insuffisance rénale chronique (MRC) et associés à de mauvais résultats globaux, mais leur évaluation n'a pas reçu l'attention méritée. Cette étude a évalué le fardeau auto-perçu par les patients sur les soignants, le bien-être psychosocial des patients atteints d'IRC et l'impact sur leur qualité de vie (QoL) dans deux hôpitaux en Nigéria.

Méthodes: Il s'agissait d'une étude transversale qui évaluait le fardeau auto-perçu des patients sur les soignants, la qualité de vie, l'anxiété et la dépression à l'aide du questionnaire Cousineau, du questionnaire SF-12 modifié et de l'échelle d'anxiété et de dépression hospitalière respectivement chez les patients atteints d'IRC et les témoins.

Résultats: Il y avait 141 participants à l'étude comprenant 50 patients hémodialysés d'entretien (HD), 41 patients atteints d'IRC en pré-dialyse et 50 témoins. Le rapport homme: femme pour les patients atteints d'IRC et le groupe témoin était respectivement de 1,1 et 1,4. La prévalence des symptômes dépressifs et anxieux chez les patients atteints d'IRC était de 46,2% et 33,0% respectivement. Le score de qualité de vie était significativement plus élevé dans le groupe témoin par rapport aux patients atteints d'IRC ($p = <0,001$). Le score de qualité de vie était significativement plus faible chez les patients HD par rapport aux patients atteints d'IRC en pré-dialyse ($p = <0,001$). Le score d'autoévaluation du fardeau était significativement plus élevé dans le groupe HD que dans le groupe avant la dialyse CKD ($p = <0,001$). Il y avait une corrélation négative significative entre la qualité de vie et l'auto-perception du fardeau, le score d'anxiété hospitalière et le score de dépression hospitalière ($p = <0,001$).

Conclusion: Les maux psychologiques sont très répandus chez les patients atteints d'IRC et ont un impact négatif sur leur qualité de vie. Une évaluation et une prise en charge régulières de ces maux doivent être intégrées dans la prise en charge des patients atteints d'IRC. **WAJM 2021; 38 (4): 335-341**

Mots clés: anxiété, dépression, qualité de vie, fardeau pour les soignants, maladie rénale chronique

¹Department of Internal Medicine, University of Medical Sciences, Ondo City, Nigeria

²Department of Internal Medicine, University of Benin Teaching Hospital, Benin City, Nigeria

*Correspondence: Dr O.S Abolarin, Email: datsy4life@hotmail.com